

CHUKA



UNIVERSITY

UNIVERSITY EXAMINATIONS

FIRST YEAR EXAMINATION FOR THE AWARD OF DEGREE OF BACHELOR OF SCIENCE AND BACHELOR OF EDUCATION (SCIENCE)

CHEM 120: PHYSICAL CHEMISTRY I

STREAMS: BSC (CHEM, MATHS, BIOL, BIOCHEM, INDUSTRIAL. CHEM, BIOMED) TIME: 2 HOURS

DAY/DATE: FRIDAY 18/12/2020

8.30 A.M. – 10.30 A.M.


INSTRUCTIONS: Answer question ONE and any other TWO questions

QUESTION ONE (30 MARKS)

- (a) (i) State the Avogadro's law (1 mark)
- (ii) State what can you deduce from the fact that at STP 22.4 dm³ of carbon dioxide contain more than 6.02×10^{23} molecules (2 marks)
- (iii) On reacting potassium chromate (VII) (potassium chlorate), KClO₄, with fluorosulphonic acid FSO₃H a gas X is evolved. A 0.245 g of X are found to occupy 112cm³ at 293 k and at pressure of 5.20×10^4 Pa. Calculate the relative molecular mass of X. { (K = 39.098, Cl = 34.45, O = 15.999 g/mol) R = $8.314 \text{ JK}^{-1}\text{mol}^{-1}$ } (2 marks)
- (iv) Describe how the syringe method can be used to determine the relative molecular mass of a gas and vapour (volatile liquids) (5 marks)
- (b) (i) Write short notes on the order of reaction and rate constants (3 marks)

CHEM 120

- (ii) The progress of the reaction below can be followed by using the fact that compound A reacts with acidified potassium iodide, liberating iodine whereas compound B does not.



A series of experiments was carried out to determine the initial rate of reaction for various concentrations of compound A. The following data were obtained.

Concentration of compound A (mol dm^{-3})	Initial rate ($\text{mol dm}^{-3}\text{s}^{-1}$)
0.060	3.12×10^{-4}
0.120	6.23×10^{-4}
0.180	9.38×10^{-4}
0.240	12.5×10^{-4}

From the data in the table, deduce the order of reaction with respect to compound A. Explain your reasoning (1 mark)

- (iii) In an investigation of the recombination of X atoms to give X_2 molecules in the gas phase in the presence of argon, that is the reaction $\text{X} + \text{X} + \text{Ar} \rightarrow \text{X}_2 + \text{Ar}$ the following data were obtained.

With the concentration of argon fixed at $1.0 \times 10^{-3} \text{ mol dm}^{-3}$

Concentration [X] (mol dm^{-3})	Initial rate $\frac{d[\text{X}_2]}{dt}$ ($\text{mol dm}^{-3}\text{s}^{-1}$)
1.0×10^{-5}	8.70×10^{-4}
2.0×10^{-5}	3.48×10^{-3}
4.0×10^{-5}	1.39×10^{-2}

With the concentration of X fixed at $1.0 \times 10^{-5} \text{ mol dm}^{-3}$

Initial concentration [Ar] (mol dm^{-3})	Initial rate $\frac{d[\text{X}_2]}{dt}$ ($\text{mol dm}^{-3}\text{s}^{-1}$)
1.0×10^{-3}	8.70×10^{-4}
5.0×10^{-3}	4.35×10^{-3}
1.0×10^{-2}	8.69×10^{-3}

Find the order of reaction with respect to [X] and [Ar], hence determine the overall velocity constant for the formation of X_2 molecules (3 marks)

CHEM 120

- (c) Draw the various types of curves showing variation of the rate with temperature (5 marks)
- (d) At certain temperature, equilibrium constant for the reaction $\text{CO} + \text{H}_2\text{O} \rightleftharpoons \text{CO}_2 + \text{H}_2$ is 5.0. If the reaction started with initial partial pressures of CO and steam as 1 atmosphere and 10 atmospheres, calculate the partial pressure of all the gases when the equilibrium state is reached. (5 marks)
- (e) A 2.5 M solution of a weak monobasic acid has the same pH as 0.017M HCl. Assuming that the HCl is completely ionized;
- calculate the pH of these solutions (1 mark)
 - determine the degree of ionization of the weak acid in its 2.5M solution (2 marks)

QUESTION TWO (20 MARKS)

- a) The rate constant for the reaction $\text{H}_2(\text{g}) + \text{I}_2(\text{g}) \rightarrow 2\text{HI}(\text{g})$ is $5.4 \times 10^{-4} \text{ m}^{-1}\text{s}^{-1}$ at 326°C . At 410°C the rate constant was found to be $2.8 \times 10^{-2} \text{ m}^{-1}\text{s}^{-1}$ calculate the
- Activation energy ($R = 8.314 \text{ JK}^{-1}\text{mol}^{-1}$) (3 marks)
 - High temperature limiting rate constant for this reaction. (2 marks)
- b) Briefly explain why the study of the effect of temperature on rates is important (3 marks)
- c) The table below gives values for the velocity constant k , of the reaction between potassium hydroxide and bromoethane in ethanol at a series of temperature, T


$K (\text{dm}^3\text{mol}^{-1}\text{s}^{-1})$	T (k)
0.182	305.0
0.466	313.0
1.35	323.1
3.31	332.7
10.2	343.6
22.6	353.0

Calculate the value for the activation energy of the reaction ($R = 8.31 \text{ JK}^{-1}\text{mol}^{-1}$)

(5 marks)

CHEM 120

d) The following data are for the reaction



[A]	[B]	Initial rate (mol litre ⁻¹ sec ⁻¹)
0.1	0.1	4.0×10^{-4}
0.2	0.2	1.6×10^{-3}
0.5	0.1	1.0×10^{-2}
0.5	0.5	1.0×10^{-2}

- (i) Determine the order with respect to A and B (4 marks)
- (ii) Calculate the rate constant (2 marks)
- (iii) Determine the reaction rate when the concentrations of A and B are 0.2M and 0.35M respectively. (1 mark)

QUESTION THREE (20 MARKS)

a) This question concerns the following reversible process



Explain the effect on the position of equilibrium on;

- (i) Increasing the pressure at constant temperature (1 marks)
- (ii) Increasing the temperature at constant pressure (1¹/₂ mark)
- (iii) Bearing in mind your answer for part (i), how do you account for the fact that industrially a temperature of about 800K, a pressure of 1 or 2 atm and a catalyst are used. (2 marks)
- (iv) At 800K, the equilibrium partial pressures in atmospheres are; $P_{\text{SO}_2} = 0.1$, $P_{\text{O}_2} = 0.7$ and $P_{\text{SO}_3} = 0.8$. Calculate K_p (1¹/₂ mark)
- b) Consider the reaction $\text{PCl}_5(\text{g}) \rightleftharpoons \text{PCl}_3(\text{g}) + \text{Cl}_2(\text{g})$. For such a system of total volume, dm³ and a total pressure of 202.6 kPa, at equilibrium the percentage dissociation of $\text{PCl}_5(\text{g})$ at temperatures of 200°C and 300°C are 48.5 and 97.0 respectively.
- (i) Calculate the value for K_p at 200°C (4 marks)

CHEM 120

- (ii) Calculate the percentage dissociation of PCl_5 that would result if the pressure of the system was reduced to 135.1 kPa. {Temperature remains constant} (2 marks)
- (iii) How is the value obtained in (ii) explained in terms of Le chatelier's principle? (1 mark)
- c) State the law of Gay-Lussac and show how it is related to Avogadro's principle (3 marks)
- d) The reaction, $2\text{N}_2\text{O}_5 \rightarrow 4\text{NO}_2 + \text{O}_2$, is forming NO_2 at the rate of 0.0076 mol/L/sec at some time. Calculate;
- (i) The rate of appearance of O_2 at this time (2 marks)
- (ii) The rate of disappearance of N_2O_5 at this time (2 marks)

QUESTION FOUR (20 MARKS)

- a) Distinguish between electrolytic cell and electrochemical cell with aid of suitable diagrams (4 marks)
- b) Will Ce^{3+} ($a=1$) ions reduce chlorine to Cl^- ($a=1$) ions at 298 K according to the reaction
- $$\text{Ce}^{3+} (a=1) + \frac{1}{2} \text{Cl}_2 (p = 1 \text{ atm}) \rightleftharpoons \text{Ce}^{4+} (a=1) + \text{Cl}^- (a=1)$$
- Given that
- $$E_{\text{Ce}^{4+}/\text{Ce}^{3+}/\text{Pt}}^{\circ} = 1.82 \text{ V}$$
- $$E_{\text{Cl}^-/\text{Cl}_2/\text{Pt}}^{\circ} = 1.3595 \text{ V} \quad (4 \text{ marks})$$
- c) Using the thermochemical data given below at 298 K,
- $$2\text{OF}_2 (\text{g}) \rightarrow \text{O}_2 (\text{g}) + 2\text{F}_2 (\text{g}) \quad \Delta H^{\circ} = -49.4 \text{ KJ}$$
- $$2\text{ClF} (\text{g}) + \text{O}_2 (\text{g}) \rightarrow \text{Cl}_2\text{O} (\text{g}) + \text{OF}_2 (\text{g}) \quad \Delta H^{\circ} = +205.6 \text{ KJ}$$
- $$\text{ClF}_3 (\text{g}) + \text{O}_2 (\text{g}) \rightarrow \frac{1}{2} \text{Cl}_2\text{O} (\text{g}) + \frac{3}{2} \text{OF}_2 (\text{g}) \quad \Delta H^{\circ} = +266.7 \text{ KJ}$$
- Calculate the change in enthalpy (ΔH°) for the following reaction
- $$\text{ClF} (\text{g}) + \text{F}_2 (\text{g}) \rightarrow \text{ClF}_3 (\text{g}) \quad \Delta H^{\circ} = ? \quad (3 \text{ marks})$$
- d) The first proton of sulphuric acid is completely ionized but the second proton is only partially dissociated with an acidity constant K_{a2} of 1.2×10^{-2} . Calculate the pH of the solution. (4 marks)

CHEM 120

- e) The hydrogen sulphate ion, HSO_4^- , is a moderately strong Bronsted acid with a K_a of 1.0×10^{-2}
- i. Write the equilibrium expression for this acid (1 mark)
 - ii. Calculate the value of $\{\text{H}^+\}$ in 0.010M HSO_4^- (Furnished by the salt, NaHSO_4). (3 marks)
 - iii. Calculate the percentage ionization of HSO_4^- into H^+ and SO_4^{2-} in 0.010M HSO_4^- (1 mark)
-